

The frequency of electron-ion collisions in a hot plasma

Fedor V.Prigara

*Institute of Microelectronics and Informatics, Russian Academy of Sciences,
21 Universitetskaya, Yaroslavl 150007, Russia**

(Dated: February 2, 2008)

Abstract

It is shown that the anomalous resistivity, thermal conductivity, and magnetic pressure of hot plasmas can be explained by the assumption that the collisional electron-ion cross-section becomes constant above some critical temperature. This constant is determined by the size of ion (its electron envelope). It is shown also that this assumption follows from the consideration of interaction of a hot plasma with thermal radiation.

PACS numbers: 52.20.Fs, 52.25.Fi, 52.25.Qs, 95.30.Qd

It is adopted normally that the only coupling between ions and electrons in hot plasmas is Coulomb collisions which is rather weak (see, e.g., [1]). The frequency of electron-ion collisions in this case is given by the formula [2, 3]

$$\nu_{ei} = nv_{Te}\sigma_{ei} \approx 2\pi e^4 n \log(n\lambda_D^3) / (m_e^2 v_{Te}^3), \quad (1)$$

where $n = n_e = n_i$ is the electron and ion density, v_{Te} is the thermal velocity of electrons, σ_{ei} is the electron-ion collisional cross-section, m_e and e are the mass and charge of electron respectively, and λ_D is the Debye length. It is clear from equation (1) that the Coulomb collisional cross-section is very small at high temperatures, $\sigma_{ei} \propto T_e^{-2}$, where T_e is the electron temperature.

Here we show that this assumption on the electron-ion collisional cross-section is not correct due to the finite dimensions of ions in a hot plasma interacting with thermal radiation.

The following effects indicate that the collisional properties of hot plasmas with T_e larger than $T_0 \cong 3keV$, where T_0 is the inversion temperature [8], essentially differ from those of cold plasmas: 1) anomalous resistivity of a plasma in strong electric fields [4, 5]; 2) the linear dependence of magnetic pressure on the magnetic field strength inferred from the pressure and magnetic field profiles for compact radio sources [6]; 3) radiation-induced solitary waves in hot accretion disks [7].

Below we consider these effects in more detail.

Consider current instabilities in a plasma in the absence of magnetic field. With respect to the voltage-current characteristics, $j = j(E)$, there are four regimes of electric fields (see also [5]):

- 1) a classical regime ($j \propto E$) for very low electric fields;
- 2) quasi-linear regime, $j \cong enc_s = const$, where c_s is the sound velocity;
- 3) the heating of a plasma, when the electron temperature, T_e , increases, and the voltage-current characteristics is $j \propto E^\alpha$;
- 4) the regime of the radiative cooling of a plasma, when $T_e \approx const$, and $j \propto E$.

The last regime is normally attributed to the Buneman instability.

In the region of $j = const$, the collisional electron-ion cross-section is $\sigma_{ei} \cong \pi (e^2/T_e)^2$, and the equation $T_e \cong eEl$, where $l = 1/n\sigma_{ei}$ is the mean free path of electrons, gives $T_e \cong \pi ne^3/E$. It means the absence of plasma heating, $T_e \approx const$. It is the region of the ion-sound instability.

Due to the turbulent heating of a plasma, the electron-ion collisional cross-section becomes a constant, σ_0 , which is determined by the atomic size (see below). In this case, which corresponds to a hot plasma in a sufficiently strong electric field, $E > E_c \cong T_e n \sigma_0 / e$, the current velocity, u_e , is proportional to the electric field, E .

In the intermediate region of electric fields, where $u_e \cong v_{Te}$, we have $m_e v_{Te}^2 \cong eE / (n \sigma_0)$, and therefore $v_{Te} \propto E^{1/2}$. Since

$$j = e^2 E / (m_e v_{Te} \sigma_0) \quad (2)$$

in this regime $j \propto E^{1/2}$. (Compare this result with the relation $j \propto E^{1/3}$ given by Galeev and Sagdeev [5]).

The critical value of the electric field, E_c , is determined by equation

$$u_e = j / en = eE / nm_e v_{Te} \sigma_0 \cong v_{Te}, \quad (3)$$

which gives

$$E_c \cong m_e v_{Te}^2 n \sigma_0 / e \cong T_e n \sigma_0 / e. \quad (4)$$

It follows from the above considerations, that in a sufficiently hot plasmas the electron-ion collisional cross-section is a constant, σ_0 , and the frequency of electron-ion collisions is given by a formula

$$\nu_{ei} = n v_{Te} \sigma_{ei} \cong n v_{Te} \sigma_0. \quad (5)$$

The cross-section σ_0 is much larger than the Coulomb collisional cross-section at these temperatures, and this determines the anomalous resistivity and thermal conductivity [3] of a plasma.

The origin of the constant cross-section σ_0 is as follows. A hot plasma with the temperature T_e larger than $T_0 \cong 3keV$ is intensely interacting with the field of thermal radiation. At temperatures $T > T_0$ the stimulated radiation processes dominate this interaction [8]. Thermal radiation induces radiative transitions in the system of electron and ion which corresponds to the transition of electron from the free to the bounded state.

Thus, in a hot plasma interacting with thermal radiation, the bounded states of electrons and ions restore, leading to the change of collisional properties of a hot plasma. In this

case, the electron-ion collisional cross-section has an order of magnitude of the atomic cross-section, $\sigma_0 \cong 10^{-15} \text{cm}^2$.

Consider now a current tube with the cross-section $S = \pi l^2$ and the length d placed in a magnetic field B . The current tube is under the action of Ampere's force:

$$F = BId/c, \quad (6)$$

where $I = jS$ is the current, j is the current density, and c is the speed of light. It is assumed that the current tube is perpendicular to the magnetic field force lines.

Suggesting that the quasilinear regime of the voltage-current characteristics is realized, we can put

$$j = enc_s, \quad (7)$$

where c_s is the sound velocity.

The area of the lateral surface of the current tube is $A = \pi ld$, and now we can calculate the pressure produced by the magnetic field on the current tube:

$$p_B = F/A = eBnlc_s/c. \quad (8)$$

Substituting l by the mean free path of electrons with respect to the scattering on the ions $l_{ei} = 1/n\sigma_0$, where σ_0 is the above high-temperature electron-ion collisional cross-section, we find that

$$p_B = eBc_s/\sigma_0c. \quad (9)$$

The equation (9) gives the linear dependence of the magnetic pressure on the magnetic field strength, contrary to the quadratic dependence given by magneto-hydrodynamics for cold plasmas (e.g., [4]). For a hot plasma interacting with thermal radiation, we can assume

$$c_s = (T_0/m_i)^{1/2}, \quad (10)$$

where T_0 is the inversion temperature, and m_i is the mass of ions.

In this case the magnetic pressure is simply proportional to the magnetic field strength, the coefficient in this relation being a constant.

Hot plasmas interacting with thermal radiation are characteristic for accretion disks around compact astrophysical objects such as active galactic nuclei and young pulsars.

In the case of flat-spectrum active galactic nuclei [8, 9, 10, 11] the density, temperature, and pressure profiles have a form $n \propto r^{-1/2}$, $T \propto r^{-1}$, $P = nkT \propto r^{-3/2}$. Since for compact radio sources $B \propto r^{-3/2}$, from equation (9) we obtain $P \propto P_B$. The latter relationship confirms equation (9). Here the density, temperature and pressure profiles are the same as those in the convection-dominated accretion flow (CDAF) models [9]. The magnetic field profile is different since the expression for the magnetic pressure is changed. The ratio of magnetic to gas pressure will be fixed similar to the CDAF models.

It is adopted normally (e.g., [1]) that, in a hot plasma of an accretion disk, the only coupling between ions and electrons is Coulomb collisions which is rather weak, and, since the radiation of electrons is much stronger than that of ions, the ion temperature, T_i , is much higher than that of electrons, T_e .

It follows from the above consideration that such an assumption is not correct. The coupling between electrons and ions in a hot plasma interacting with thermal radiation is strong, and therefore such a hot plasma is roughly the one-temperature plasma, i.e. $T_e \approx T_i$. This conclusion is very important for the models of accretion flows in compact astrophysical objects. In particular, it puts under the question the explanation of the phenomenon of low-luminosity active galactic nuclei by the models of accretion flows based on the assumption that T_i is much higher than T_e (see, e.g., [1, 9, 10]). It is not surprising that these models encounter difficulties in the explanation of observational properties of low-luminosity active galactic nuclei.

Another field where the strong coupling between electrons and ions in a hot plasma can essentially change the situation is the theory of envelope (Langmuir) solitary waves [12]. In the last theory the interaction between Langmuir and ion-sound waves play an important role.

-
- [1] F.Yuan, and A.Zdziarski, *Luminous hot accretion flows: the origin of X-ray emission of Seyfert galaxies and black hole binaries*, Month. Not. R. Astron. Soc. (submitted), E-print archives, astro-ph/0401058 (2004).
- [2] F.F.Chen, *Introduction to Plasma Physics and Controlled Fusion, Vol. 1: Plasma Physics* (Plenum Press, New York, 1984).
- [3] V.P.Silin, Usp. Fiz. Nauk, **172**, 1021 (2002), Physics- Uspekhi, **45**, 955 (2002).
- [4] B.B.Kadomtsev, *Collective Phenomena in Plasmas* (Nauka, Moscow, 1988).
- [5] A.A.Galeev and R.Z.Sagdeev, in *Handbook of Plasma Physics*, eds. M.N.Rosenbluth and R.Z.Sagdeev, Vol. 2: *Basic Plasma Physics II*, ed. by A.A.Galeev and R.N.Sudan (North- Holland Physics Publishers, Amsterdam, 1984).
- [6] F.V.Prigara, in *The Magnetized Interstellar Medium*, 8-12 September 2003, Antalya, Turkey, astro-ph/0304243.
- [7] F.V.Prigara, *Radiation-induced solitary waves in hot plasmas*, Phys. Plasmas (submitted), physics/0404087.
- [8] F.V.Prigara, in *Gamow Memorial International Conference*, 8-14 August 2004, Odessa, Ukraine, astro-ph/0310491.
- [9] N.M.Nagar, A.S.Wilson, and H.Falcke, Astrophys. J., **559**, L87 (2001).
- [10] J.S.Ulvestad and L.C.Ho, Astrophys. J., **562**, L133 (2001).
- [11] F.V.Prigara, *Astron. Nachr.*, **324**, No. S1, 425 (2003).
- [12] V.E.Zakharov, in *Handbook of Plasma Physics*, eds. M.N.Rosenbluth and R.Z.Sagdeev, Vol. 2: *Basic Plasma Physics II*, ed. by A.A.Galeev and R.N.Sudan (North-Holland Physics Publishers, Amsterdam, 1984).

* Electronic address: fprigara@imras.yar.ru